

Negative Pressure Wound Therapy Referral Information Sheet

Before initiation of Negative Pressure Wound Therapy (NPWT), the ordering physician / Wound Care Clinician must complete the following information

Date: _____ Address: _____

Client Name: _____

BRN: _____

Date of Birth (d/m/y): _____

Wound History: Diagnosis (Check one):



- Pilonidal sinuses Large surgical wound Mediastinal wound Necrotizing fasciitis wounds
 Pressure ulcers Orthopaedic wound Traumatic wounds Diabetic foot

Age of wound: _____

What advanced-wound Rx has been previously used to treat this wound?

What has the response been? wound static wound improving

Wound measurements & description: Length: _____ cm x width: _____ cm x Depth: _____ cm



Expected therapy goals: (i.e. Flap/Graft/Closure/Prep for Surgery) _____ in _____ weeks.

Without NPWT Therapy, how long would this/these wound(s) take to heal (approx.) _____ weeks

Indicate what the frequency would be for conventional dressing changes:

- OD BID TID 3 x week Other _____

Current Pain Scale: 1 - 2- 3 -4 -5 -6 -7 -8 -9 -10 _____

Analgesia: _____

Wounds must have the following criteria to be eligible for NPWT therapy:

- Once a day (OD), twice a day (BID), or three times a day (TID) dressing changes.
- Acute wounds/traumatic wounds.
- Surgical wounds.
- Stage III to IV Pressure Ulcers of recent occurrence.
- Dehisced wounds.
- Diabetic foot ulcers following surgical procedures or sharp debridement with viable tissue (where ABPI > 0.5.)
- Arterial insufficiency (where ABPI < 0.5.)

No NPWT therapy contraindications exist:

- Presence of necrotic tissue. (Wound must be debrided so that at least 80% of base contains healthy granulation tissue.)
- Nutritional status is not adequate to support healing. (e.g. Braden nutritional score < 3, Nutritional compromise with serum albumin < 35 g/dl, or pre-albumin level < 16 mg/dL.)
- Severe excoriation or periwound.
- An unexplored fistula to organs or body cavities (other than chronic enteric fistulas.)
- Unresolved, untreated osteomyelitis and any infection that is untreated prior to application.
- Malignancy or cancer in wound margins.
- Unresolved bleeding following debridement. Exposed blood vessels and/or organs.
- Client experiencing difficult homeostasis after debridement.



Negative Pressure Wound Therapy Referral Information Sheet

Precautions have been addressed:

- Sharp fragments of bone must be removed prior to initiation of NPWT.
- Exposed tendons, ligaments and nerves must be covered with meshed non-adherent material before the NPWT dressing is applied.
- Client receiving anticoagulants with stable INRs.
- Not experiencing active bleeding or anemia.
- Immunodeficient disease, for example Leukemia, HIV.
- Haematologic disorders well-controlled.
- Diabetes or hypertension well-controlled.
- No current abuse of drugs or alcohol.
- Systemic steroids.
- If the location of the wound interferes with the therapy by preventing a sustainable seal of the drape, the NPWT will be discontinued.

Discontinuation Criteria:

- When there is no measurable progress to wound healing within two weeks;
- When there is not 20-40 percent reduction in the size of the wound within three to four weeks;
- The wound has healed such that the foam no longer fits the wound;
- The goals for healing have been met;
- If any of the following occur: bleeding, bruising, unmanaged pain in response to the therapy, an occlusive seal cannot be achieved, the client does not comply with the treatment regime, or the wound deteriorates.
- Regardless of decrease in size, if the wound is healing as expected the NPWT will be discontinued by the end of 6 to 8 weeks of treatment

The Physician has assessed that NPWT is safe to use for this client: YES NO

NPWT TREATMENT PLAN – Identify treatment type, dressing type, size, and delivery required:

- KCI ActiVAC** Granufoam Kit: Small Medium Large
 Whitefoam Kit: Small Large
 300 ml Cannister

- Nanova Therapy Unit** Nanova Small Dressing Nanova Medium Dressing Nanova Large Dressing

DELIVERY: Regular Next-Day Home Delivery Delivery date required: _____

Initial Settings ActiVAC: (1st 48 hours all wounds) Intermittent (if wound appropriate, after 48 hrs)

- 25mm/Hg 50mm/Hg 75mm/Hg 100mm/Hg 125mm/Hg 150mm/Hg 175mm/Hg
 200mm/Hg

Provide alternate moist wound dressing treatment should the NPWT needs to be interrupted or discontinued:

At Nurse's discretion or Primary dressing: _____ Secondary dressing _____ Dressing change frequency: _____

Name of Institution _____

Physician/Wound Specialist: _____

Signature: _____

For LHIN Use Only:

SEND completed form to Vendor Yurek's with Service Referral –Pharmacy Consultation - via HPG.

Yureks Phone Number: 1-888-631-6502 **COPY TO:** Manager, Client Services

Thank you for your time and consideration.

